

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A01N 57/20, 25/30, C11D 1/94, 3/43 // 1/06	A1	(11) International Publication Number: WO 00/38523 (43) International Publication Date: 6 July 2000 (06.07.00)
(21) International Application Number: PCT/EP99/10269 (22) International Filing Date: 22 December 1999 (22.12.99) (30) Priority Data: 9828258.5 23 December 1998 (23.12.98) GB 9901205.6 21 January 1999 (21.01.99) GB (71) Applicant (for all designated States except US): ALBRIGHT & WILSON UK LIMITED [GB/GB]; 210-222 Hagley Road West, Oldbury, West Midlands B68 0NN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): MILLE, Fabien, Hervé, Joseph [FR/FR]; 1, Porte au Rupt, F-44200 Commercy (FR). OXFORD, Philip, James [GB/GB]; 82 Sterling Place, South Ealing, London W5 4RB (GB). (74) Agent: SAVIDGE, Roger, Gordon, Madgwick; Albright & Wilson UK Limited, 210-222 Hagley Road West, Oldbury, West Midlands B68 0NN (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: HERBICIDAL COMPOSITIONS AND SURFACTANT CONCENTRATES (57) Abstract <p>A herbicidal solution contains from 30 % by weight to saturation of a water soluble glyphosate salt and from 8 to 20 % by weight of surfactant comprising from 10 to 100 % by weight based on the total weight of surfactant of an amphoteric surfactant and from 0 to 90 % by weight based on the total weight of surfactant of ether carboxylate, said solution containing less than 0.035 % by weight of sodium ion. A pre-blended surfactant concentrate contains from 10 to 70 % by weight ether carboxylate, from 20 to 80 % by weight of low salt amphoteric surfactant and from 10 to 50 % by weight of solvent.</p> <div style="text-align: right; margin-top: 20px;"><p>= EP 1139761 B1</p><p>= US 5500784</p><p>= US 6653257</p></div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

HERBICIDAL COMPOSITIONS AND SURFACTANT CONCENTRATES

The present invention relates to concentrated herbicidal compositions comprising water soluble glyphosate salts and a surfactant.

A number of formulations have been proposed whereby the herbicide N-(phosphonomethyl)glycine, alias glyphosate, may be supplied as a concentrated aqueous solution with a surfactant synergist which aids wetting and penetration, when the composition is diluted and applied to herbage.

Factors governing the choice of surfactant include wetting power, herbicidal or synergistic action, environmental profile and ability to form stable solutions with glyphosate at as high a concentration as possible, as well as cost.

The surfactants which have so far proved most cost effective for these purposes have been ethoxylated amines. The latter however have a poor environmental profile being biotoxic and poorly biodegradable. There is a demand for a more environmentally acceptable alternative to amine ethoxylates.

We have discovered that ether carboxylates meet most of the above criteria but are insufficiently soluble in concentrated glyphosate solutions. We have further noted that amphoteric surfactants also meet most of the above criteria and can act as cosurfactants, solubilising ether carboxylates, but solutions of commercial amphoteric surfactants in concentrated glyphosate are unstable and have been found to deposit crystals of glyphosate salts.

Amphoteric surfactants comprise betaines, sulpho betaines and phosphobetaines. However the great majority of the amphoteric surfactants sold commercially are betaines. The betaines of commerce are normally made by quaternising an amine by reaction with sodium chloracetate. This reaction forms sodium chloride as an unavoidable by-product. As a result commercial betaines contain about 6 to 12% of sodium chloride. We have now discovered that it is possible to use amphoteric surfactants at the required levels in glyphosate solution concentrates without instability, if the sodium chloride content is substantially reduced. Low salt betaines typically contain less than 3% preferably less than 2%, more preferably less than 1% e.g. less than 0.5% especially less than 0.2% sodium ion expressed as wt. sodium chloride based on the weight of surfactant.


The invention provides a herbicidal aqueous solution comprising from 30% by weight, to saturation of a water soluble glyphosate salt and from 8 to 20% by weight of surfactant comprising 10 to 100% by weight based on the total weight of surfactant of an amphoteric surfactant and from 0 to 90% by weight based on the total weight of surfactant of ether carboxylate, said solution containing less than 0.035% by weight of sodium ion.

Sodium ion may be removed from amphoteric surfactants, either by electrosmosis, e.g. as described in our GB 1 525 692 or in EP O 736 521, or by membrane filtration, for example as described in EP O 626 881, or, less preferably, by displacing sodium ion with, for example, potassium or ammonium, e.g. using ion exchange. Alternatively, it is possible to prepare betaines with low salt levels by quaternising with acrylic acid.

The glyphosate is preferably present as its potassium, ammonium, C_{2 to 3} amine or mono or di ethanolamine salt, or metho sulphate or as a mixture of two or more of said salts. Particularly preferred is the isopropylamine salt.

We prefer that the total level of inorganic salt be less than 0.39% by weight.

The amphoteric surfactant is preferably a betaine, e.g. a betaine of the formula: $RR^1_2N^+CH_2COO^-$, wherein R is an alkyl, alkenyl or alkyl phenyl group having an average of from 6 to 20, e.g. 8 to 14 aliphatic carbon atoms and R^1 is an alkyl or hydroxy alkyl group having an average of from 1 to 4 carbon atoms. Particularly preferred are the so called quaternary imidazoline betaines, also called amphoteric acetates, and commonly ascribed the formula:



(although they are actually present predominantly as the corresponding non cyclic amido amine) wherein R and R' are alkyl, alkenyl, cycloalkyl, alkaryl or hydroxyalkyl groups having an average of from 1 to 20 aliphatic carbon atoms and R preferably has an average of from 6 to 20, e.g. 8 to 14 aliphatic carbon atoms and R' preferably has 1 to 4 carbon atoms. Other amphoteric surfactants for use according to our invention include alkyl amine polyalkoxy sulphates, sulphobetaines and other quaternary amine or quaternised imidazoline sulphonic acids and their salts, and Zwitterionic surfactants, e.g. N-alkyl taurines, carboxylated amido amines such as $RCONH(CH_2)_nN^+R'_2CH_2COO^-$ where n is 2 to 4, and amino acids having, in each case, hydrocarbon groups capable of conferring surfactant properties (e.g. alkyl, cycloalkyl alkenyl or alkaryl groups having from 6 to 20 aliphatic carbon atoms). Typical examples include 2-tallow alkyl, 1-tallow amido alkyl, 1-carboxymethyl imidazoline, 2-coconut alkyl N-carboxymethyl 2 (hydroxyalkyl) imidazoline, alkyl amido ethyl, propyl or butyl dimethyl betaine and C_{6-20} alkyl dimethyl betaine. Generally speaking any water soluble amphoteric or Zwitterionic surfactant compound which comprises a hydrophobic portion including C_{6-20} alkyl or alkenyl group and a hydrophilic portion containing an amine or quaternary ammonium group and a carboxylate, sulphate or sulphonic acid group may be used in our invention.

We particularly prefer short chain alkyl amido betaines, amphoteric acetates and alkyl betaines in which the alkyl group has 6 to 10 carbon atoms since they give very low foaming and permit high levels of ether carboxylate to be solubilised.

The surfactant preferably comprises at least 20% more preferably at least 30%, most preferably at least 40% e.g. 50 to 100% by weight of the amphoteric surfactant based on the total weight of surfactant.

We strongly prefer that the surfactant comprises an ether carboxylate. The ether carboxylate is preferably present in an amount of at least 10%, preferably more than 20% e.g. 30 to 60% by weight based on the total weight of surfactant. Generally the shorter the long alkyl chain of the betaines the higher the proportion of ether carboxylate that can be readily dissolved in a stable solution.

The ether carboxylate preferably has the formula $R(OCH_2CH_2)_n OCH_2CO_2^-$ wherein R is a straight or branched alkyl, alkenyl, alkylphenyl or polypropylene oxy group having from 6 to 20 e.g. 8 to 14 aliphatic carbon atoms and n is from 1 to 30, preferably, 2 to 20 e.g. 3 to 10.

The counter ion of the ether carboxylate may preferably comprise ammonium, potassium and or an amine or alkanolamine having up to six carbon atoms.


The surfactant may optionally but preferably comprise up to 20% by weight, e.g. 5 to 10% based on the total weight of surfactant, of a non-ionic wetting agent. The non-ionic wetting agent is preferably an ethoxylated alcohol such as a C_6 to 25 straight or branched alkyl 1 to 30 mole ethoxylate e.g. a C_{10} to 18 alkyl 5 to 20 mole ethoxylate. The wetting agent gives better coverage of the leaf and superior herbicidal action.

We prefer that the surfactants should be pre-blended as a concentrate for supply to formulators of glyphosate compositions. To maintain sufficient mobility and

to formulators of glyphosate compositions. To maintain sufficient mobility and dispersibility of such concentrates we prefer to add a suitable water miscible solvent such as a water miscible glycol or glycol ether, e.g. polyethylene glycol having a mean mol weight between 90 and 600 especially 130 to 250.

Such concentrates containing from 10 to 70% by weight ether carboxylate, from 20 to 80% by weight of low salt amphoteric surfactant and from 10 to 50% by weight of solvent are a further embodiment of the invention.

Preferably said concentrates additionally contain from 3 to 20% of a non-ionic wetting agent and/or from 1 to 6% of a sequestrant such as an aminophosphonate, e.g. of the formula.



where n is from 0 to 10 preferably 2 to 6 and M is a compatible cation to provide a water soluble product.

The formulation may typically contain from 1 to 3% of sequestrant.

The concentrates contain, typically less than 70% water, preferably less than 60% water most preferably 10 to 55% water by weight of the concentrate.

The invention will be illustrated by the following examples :-

Example 1

A composition was prepared comprising 360g l⁻¹ isopropylamine salt of glyphosate as acid equivalent and 105g l⁻¹ active ingredient of C₁₂₋₁₄ alkyl betaine, as a 37% solution desalinated by electrodialysis to a sodium chloride level of less than 1%. The composition was a clear solution which was stable on storage in a cold incubator at 0°C and at ambient temperature.

For comparison a composition was prepared as described above, but using as surfactant a commercial C₁₂₋₁₄ alkyl betaine sold under the Registered Trade Mark "EMPIGEN" BB and containing 6% sodium chloride. The comparative example was unstable and deposited crystals of glyphosate when stored in a cold incubator at 0°C and after 2 to 3 weeks at ambient temperature.

Example 2

The formulation of claim 1 was prepared using a desalinated coco imidazoline betaine and for comparison a commercial coco imidazoline betaine sold as a 38% solids aqueous solution containing 11% sodium chloride under the Registered Trade Mark "EMPIGEN" CDR60.

The former composition was stable on storage at 0°C and at 54°C. The comparative example deposited crystals of glyphosate at both temperatures.

Example 3

The formulation of example 1 was repeated using a desalinated coco amidopropyl betaine, and, as comparison a commercial equivalent sold as "EMPIGEN" BS containing 6.5% sodium chloride. The example was stable on storage at 0°C, but the comparative example deposited crystals of glyphosate at these temperatures.

Examples 4 to 16

The following compositions containing alkyl polyethoxymethyl carboxylates of the formula $R(OCH_2CH_2)_nOCH_2CO_2H$ where R and n are as defined below and the desalinated alkyl betaine of claim 1 were all stable at 0°C for 1 week, and at 54°C for two weeks and were fully soluble in water containing 500ppm hardness.

The ether carboxylates were sold under the Registered Trade Mark "EMPICOL" and are identified as follows:-

"EMPICOL" CYJ	R = hexyl, n = 10
"EMPICOL" CVE	R = capryl, n = 6
"EMPICOL" CVH	R = capryl, n = 8
"EMPICOL" CLI	R = oleyl, n = 10
"EMPICOL" CED5	R = lauryl, n = 6

Example 4

Glyphosate isopropylamine salt:	360g/l <i>as acid equivalent</i>
Polyethylene glycol (MW200)	40g/l
Desalinated alkyl betaine	52.5g/l <i>as active matter</i>
"EMPICOL" CYJ	52.5g/l <i>as active matter</i>
Ammonia	to pH 5 +/- 0.1
Demineralised water	to 1l or 1176g

The PEG betaine and ether carboxylate were mixed to form a stable concentrate. The latter was dispersed readily in the water with the glyphosate and ammonia.

Example 5

As example 4 but replacing "EMPICOL" CYJ with "EMPICOL" CVE

Example 6

As example 4 but replacing ""EMPICOL"" CYJ with "EMPICOL" CVH

Example 7

As example 4 but replacing "EMPICOL" CYJ with "EMPICOL" CLI

Example 8

Glyphosate isopropylamine salt:	360g/l <i>as acid equivalent</i>
Polyethylene glycol	40g/l
Desalinated alkyl betaine	75g/l <i>as active matter</i>
"EMPICOL" CYJ	75g/l <i>as active matter</i>
Ammonia	to pH 5 +/- 0.1
Demineralised water	to 1l or 1176g

The PEG, ether carboxylate and betaine were mixed to form a stable concentrate which was easily dispersed in water with the glyphosate and ammonia.

Example 9

As for example 8 replacing "EMPICOL" CYJ for "EMPICOL" CVE

Example 10

As for example 8 replacing "EMPICOL" CYJ for "EMPICOL" CVH

Example 11

Glyphosate isopropylamine salt:	360g/l <i>as acid equivalent</i>
Polyethylene glycol	40g/l
Desalinated alkyl betaine	42g/l <i>as active matter</i>
"EMPICOL" CYJ	63g/l <i>as active matter</i>
Ammonia	to pH 5 +/- 0.1
Demineralised water	to 1l or 1176g

The PEG, betaine and ether carboxylate were mixed to form a stable concentrate which was easily dispersed in water with the glyphosate and ammonia.

Example 12

As for example 11, replacing "EMPICOL" CYJ for "EMPICOL" CVE

Example 13.

As for example 11 replacing "EMPICOL" CYJ by "EMPICOL" CVH

Example 14

Glyphosate isopropylamine salt:	360g/l <i>as acid equivalent</i>
Polyethylene glycol (MW200)	40g/l
Desalinated alkyl betaine	60g/l <i>as active matter</i>
"EMPICOL" CVE	90g/l <i>as active matter</i>
Ammonia	to pH 5 +/- 0.1
Demineralised water	to 11 or 1176g

Example 15

Glyphosate isopropylamine salt:	360g/l <i>as acid equivalent</i>
Polyethylene glycol (MW200)	40g/l
Desalinated alkyl betaine	60g/l <i>as active matter</i>
"EMPICOL" CVH	90g/l <i>as active matter</i>
Ammonia	to pH 5 +/- 0.1
Demineralised water	to 11 or 1176g

Example 16

Glyphosate isopropylamine salt:	360g/l <i>as acid equivalent</i>
Polyethylene glycol (MW200)	40g/l
Desalinated cocoamidopropyl betaine of example 3	52.5g/l <i>as active matter</i>
"EMPICOL" CVE	52.5g/l <i>as active matter</i>
Ammonia	to pH 5 +/- 0.1
Demineralised water	to 11 or 1176g

Examples 17 to 28

Examples 4 to 15 were repeated using a C₈ alkyl amidopropyl betaine. Stable concentrates and solutions exhibiting very low foaming were obtained.

Examples 29 to 53

Examples 4 to 28 were repeated with the addition of 6% by weight, based on the total weight of surfactant, of a fatty alcohol 7 mole ethoxylate sold commercially under the registered trademark "EMPILAN" KI7. The concentrates and solutions were all stable and the solutions exhibited improved wetting agent of leaf surfaces and herbicidal activity.

Examples 54 to 103

Examples 4 to 53 were repeated with the addition of 2% based on the weight of surfactant of a chelating agent, diethylene triamine pentakis (methylene phosphonic) acid sold commercially under the Registered trademark "BRIQUEST" 543.45AS. The concentrates were stable and the solutions gave improved tolerance of hardwater.

CLAIMS

1. A herbicidal aqueous solution comprising from 30% by weight of said solution to saturation of a water soluble glyphosate salt and from 8 to 20 by weight of said solution of surfactant, said surfactant comprising from 10 to 100% by weight based on the total weight of said surfactant of an amphoteric surfactant and from 0 to 90% by weight based on the total weight of surfactant of ether carboxylate, said solution containing less than 0.035% of sodium ion.
2. A solution according to claim 1 wherein said glyphosate is present as its potassium, ammonium, C_2 to C_3 amine, mono or diethanolamine or methosulphate salt.
3. A solution according to either of claims 1 and 2 wherein said amphoteric surfactant is betaine of the formula $RR^1_2N^+CH_2COO^-$ wherein R is an alkyl, alkenyl or alkyl phenyl group having 6 to 20 e.g. 8 to 14 aliphatic carbon atoms and each R^1 is an alkyl or hydroxyalkyl group having from 1 to 4 carbon atoms.
4. A solution according to either of claims 1 and 2 wherein said amphoteric surfactant has the formula $RCONH(CH_2)_nN^+R^1_2CH_2CO_2^-$ where R and R^1 have the same significance as in claim 3 and n is an integer from 2 to 4.
5. A solution according to either of claims 3 and 4 wherein R is an alkyl group having from 6 to 10 carbon atoms.
6. A solution according to any foregoing claim wherein said amphoteric surfactant constitutes at least 30% of the total weight of the surfactant.
7. A solution according to any foregoing claim comprising at least 30% by weight of ether carboxylate, based on the total weight of surfactant.

8. A solution according to any foregoing claim wherein said surfactant comprises up to 20% by weight thereof of a non-ionic wetting agent.
9. A solution according to any foregoing claim comprising a sequestrant.
10. A solution according to claim 9 wherein said sequestrant is an amino phosphonate of the formula $(M_2PO_3CH_2)_2N(CH_2CH_2NPO_3M_2)_nCH_2PO_3M_2$ wherein n is from 0 to 10 and M is cation such that the phosphonate is water soluble.
11. A concentrate for use in preparing solutions according to any foregoing claim comprising from 10 to 70% by weight ether carboxylate from 20 to 80% by weight of an amphoteric surfactant which contains less than 3% of sodium chloride and from 10 to 50% by weight of a water miscible solvent.
12. A concentrate according to Claim 11 containing from 3 to 20% of a non-ionic wetting agent.
13. A concentrate according to either of claims 11 and 12 wherein said solvent is a polyethylene glycol having a mean molecular weight of between 90 and 600.
14. A concentrate according to any of claim 11 to 13 containing 1 to 6% of a sequestrant.
15. An aqueous solution according to claim 1 comprising water, glyphosate and a concentrate according to any of claims 11 to 14.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/10269

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01N57/20 A01N25/30 C11D1/94 C11D3/43 //C11D1/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 498 785 A (MONSANTO EUROPE SA) 12 August 1992 (1992-08-12) page 2, line 1 - line 21 page 2, line 34 -page 3, line 7	1-15
A	EP 0 736 521 A (RHONE POULENC CHIMIE) 9 October 1996 (1996-10-09) cited in the application page 2 -page 3, line 4 page 7, line 16 - line 38 page 7, line 49 -page 8, line 19	1-15
A	US 4 243 549 A (MESSENGER EDWARD T ET AL) 6 January 1981 (1981-01-06) column 2, line 44 - line 60 column 4, line 15 - line 46 column 4, line 51 -column 5, line 65 -/-	11-14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

12 May 2000

Date of mailing of the international search report

13/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Muellners, W

INTERNATIONAL SEARCH REPORT

Intern 1al Application No
PCT/EP 99/10269

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 110, no. 24, 12 June 1989 (1989-06-12) Columbus, Ohio, US; abstract no. 215257, OZASA, YUTAKA ET AL: "Liquid cleaning compositions containing amphoteric surfactants purified by reverse osmosis" XP002137551 abstract & JP 10 183176 A (KAWAKEN FINE CHEMICALS CO. LTD.) 14 July 1998 (1998-07-14)	11-14
A	-& PATENT ABSTRACTS OF JAPAN vol. 1998, no. 12, 31 October 1998 (1998-10-31) & JP 10 183176 A (KAWAKEN FINE CHEM CO LTD.), 14 July 1998 (1998-07-14) abstract	11-14
A	WO 97 06230 A (RECKITT & COLMAN INC ;RICHTER ALAN FRANCIS (US); TARASCHI FREDERIC) 20 February 1997 (1997-02-20) page 2, line 16 - line 30 page 7, line 1 - line 21 page 8, line 21 -page 9, line 17; examples 2A,,3A,4B,,6B-9B	11-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/10269

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0498785	A	12-08-1992	AT 161144 T	15-01-1998
			AU 650091 B	09-06-1994
			AU 1060092 A	13-08-1992
			CA 2060870 A	09-08-1992
			DE 69223529 D	29-01-1998
			DE 69223529 T	02-07-1998
			DK 498785 T	09-02-1998
			ES 2112308 T	01-04-1998
			GR 3026188 T	29-05-1998
			IL 100873 A	18-02-1997
			JP 4352704 A	07-12-1992
			KR 9509173 B	16-08-1995
			NZ 241500 A	26-10-1994
			PT 100103 A,B	31-05-1993
			ZA 9200768 A	25-11-1992
EP 0736521	A	09-10-1996	FR 2732689 A	11-10-1996
			JP 8323156 A	10-12-1996
US 4243549	A	06-01-1981	AU 523665 B	12-08-1982
			AU 3832278 A	31-01-1980
			DE 2832814 A	15-02-1979
			FR 2398797 A	23-02-1979
			IT 1160447 B	11-03-1987
			JP 1607777 C	13-06-1991
			JP 54048691 A	17-04-1979
			JP 61038239 B	28-08-1986
			GB 2026437 A	06-02-1980
JP 10183176	A	14-07-1998	JP 2932377 B	09-08-1999
WO 9706230	A	20-02-1997	GB 2304111 A	12-03-1997
			GB 2304112 A	12-03-1997
			AU 700516 B	07-01-1999
			AU 6687096 A	05-03-1997
			BR 9609984 A	12-01-1999
			CA 2228427 A	20-02-1997
			CN 1198772 A	11-11-1998
			EP 0842248 A	20-05-1998
			GB 2304115 A,B	12-03-1997
			US 5591708 A	07-01-1997
			US 5728672 A	17-03-1998
			GB 2304728 A,B	26-03-1997
			US 5629280 A	13-05-1997
			US 5939374 A	17-08-1999